

THE UNITED STATES OF AMERICA

TO AMETO WHOM THESE PRESENTS SHAME COME:

Pure Seed Testing, Inc.

HELENS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH GASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE-SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HERS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC PARSHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE SAID TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR UTING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE POSES, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

FESCUE, RED

'Inverness'

In Testimone Mexicol. I have hereunto set my hand and caused the seal of the Hunt United Protection Office to be affixed at the City of Washington, D.C. this thirtieth day of January, the year two thousand and eight.

Attest

Derz-

Commissioner Plant Variety Protection Office Agricultural Marketing Service Secretary of Lure

Crystal Rose-Fricker

CAPACITY OR TITLE

President

Joseph K

Taxonomist/Plant

(See reverse for instructions and information collection burden state

Ρh . D

Breeder

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVP application form signed by the owner; (2) completed exhibits A. B. C. F. (3) for a cold control of the pvP application form signed by the owner; (2) completed exhibits A. B. C. F. (3) for a cold control of the pvP application form signed by the owner; (2) completed exhibits A. B. C. F. (3) for a cold control of the pvP application form signed by the owner; (2) completed exhibits A. B. C. F. (3) for a cold control of the pvP application form signed by the owner; (2) completed exhibits A. B. C. F. (3) for a cold control of the pvP application form signed by the owner; (2) completed exhibits A. B. C. F. (3) for a cold control of the pvP application form signed by the owner; (2) completed exhibits A. B. C. F. (3) for a cold control of the pvP application form signed by the owner; (2) completed exhibits A. B. C. F. (3) for a cold control of the pvP application form signed by the owner; (2) completed exhibits A. B. C. F. (3) for a cold control of the pvP application form signed by the owner; (2) completed exhibits A. B. C. F. (3) for a cold control of the pvP application form signed by the owner; (2) control of the pvP application form signed by the owner; (2) control of the pvP application form signed by the owner; (2) control of the pvP application form signed by the owner; (3) control of the pvP application form signed by the owner; (4) control of the pvP application form signed by the owner; (4) control of the pvP application form signed by the owner; (5) control of the pvP application form signed by the owner; (5) control of the pvP application form signed by the owner; (6) control of the pvP application form signed by the owner; (6) control of the pvP application form signed by the owner; (7) control of the pvP application form signed by the owner; (8) control of the pvP application form signed by the owner; (8) control of the pvP application form signed by the owner; (8) cont application form signed by the owner; (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$3,652 (\$432 riviir reproduce an entire planty tissue cuttine will be deposited and intalitation and reproduce and entire planty tissue cuttine will be deposited and interest and \$3,220 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfilled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 401, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$432 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

Plant Variety Protection Office Telephone: (301) 504-5518 FAX: (301) 504-5291

Homepage: http://www.ams.usda.gov/science/pvpo/pvp.htm

To avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority and provide evidence that name has been cleared by the appropriate recognized authority before the Certificate of Protection is issued. For example, for agricultural and vegetable crops, contact: Seed Branch, AMS, USDA, 10301 Baltimore Avenue, Suite 401 NAL Building, Beltsville, MD 20705. Telephone: (301) 504-5682 http://www.ams.usda.gov/lsg/seed.htm.

ITEM

- 19a. Give:
- (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- (2) the details of subsequent stages of selection and multiplication;
- evidence of uniformity and stability; and
- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 19b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 19c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 19d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 19e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 20. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 23. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 24. See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.
- 22. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)
- 23. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

The first commercial seed of this variety was sold 24 February 2003.

24, CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. The fees for filling a change of address; owner's representative; ownership or assignment; or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 1.4 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, merital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

Amended Exhibit A.

Origin and Breeding History of Inverness (PST-47TCR) Spreading Red Fescue

Inverness (PST-47TCR) strong creeping red fescue (*Festuca rubra* L. ssp. rubra) is an advanced generation synthetic resulting from a salt screening of the progeny of PST-4VB, which traces back maternally to the cultivar Florentine, and and then top crossed with plants that trace to 'Shademaster II', 'Shademaster', 'Badger', 'Flyer II' and collections from Rose City Cemetery, Portland, OR.

Summer of 1993. Fifty plants of PST-4VB (along with plants that trace to 'Shademaster II', 'Shademaster', 'Badger', 'Flyer II' and collections from Rose City Cemetery, Portland, OR) were screened in a salt bath at 6,000 mg L⁻¹. Thirteen plants of PST-4VB survived and were planted into a nursery in the fall of 1993. Other spreading red fescue salt survivors from this screening were also planted in this nursery: 'Shademaster II' (9 plants), 'Shademaster' (6 plants), 'Badger' (8 plants), 'Flyer II' (7 plants) and collections from Rose City Cemetery, Portland, OR (20 plants).

Summer 1995. All the plants were allowed to interpollinate and the following plants were harvested and designated as PST-46T:

PST-4VB (3 plants)

'Badger' (3 plants)

'Shademaster' (2 plants)

Rose City Collection (3 plants)

'Shadow' (1 plant)

'Shademaster II' (3 plants)

'Camilla' (1 plants)

Each plant was harvested separately.

Fall 1995. Seed was started from each plant and used to establish a progeny nursery of 61 plants per maternal parent. Seed was also used to establish a turf trial.

Summer 1996. The nursery was rogued for maturity, freedom of disease and color. Nine plants were harvested from the PST-4VB source and designated PST-46TCR.

Fall 1996. A 2,050-plant nursery planted by maternal lines of PST-46TCR, was established. The nursery was rogued for maturity, freedom of disease and uniformity. A total of 188 plants were harvested as breeder seed and designated as PST-47TCR the summer of 1997.

Breeder seed was also harvested from the same plants in 1998 and 1999.

Seed propagation of Inverness is limited to three generations of increase from Breeder seed: one each of Foundation, Registered and Certified. Pure Seed Testing, Inc. maintains Breeder seed in Oregon.

Inverness is a stable and uniform variety. No off-type or variants have been observed in the production or multiplication of this variety. This has been true for three years of production; breeder seed, foundation and certified.

Breeder, foundation, registered and certified seed of Inverness will produce turf and seed fields of equal quality, acceptable uniformity and good stability.

Amended Exhibit B.

Novelty Statement for Inverness (PST-47TCR, PST-47TCL) Strong Creeping Red Fescue

Inverness is most similar to Florentine, but close examination shows the following differences.

- Inverness has a heading date 4 days earlier than Florentine (Tables 1 and 1A).
- Inverness has a plant height at least 4 cm taller than Florentine (Tables 2 and 2A).

The differences in plant height for Florentine could be that Table 2 data is taken on two year old plants and Table 2A is on baby or first year measurements which are usually smaller with a fall planting date. When we looked at the raw data, reps 2 and 3 were consistently shorter than rep 1 as well so there may have been some environmental differences between reps. At this time we don't have any additional plant height data to show a significant difference between these varieties.

3. The anther color of Inverness is yellow green where Florentine is purplish.

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this collection of information is (0581-0055). The time required to complete this information collection is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact the USDA's TARGET Center at 202-720-2600 (voice and TDD). To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

> U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY PROGRAM PLANT VARIETY PROTECTION OFFICE BELTSVILLE, MD 20705

EXHIBIT C (FINE LEAVED FESCUES)

OBJECTIVE DESCRIPTION OF VARIETY FINE LEAVED FESCUES

(Festuca spp.)

200400123

NAME OF APP	LICANT(S)	TI	EMPORARY	DESIGNATION	VARIETY NAME				
Pure Seed Tes	sting, Inc.	1	PST-47T	CR, PST-47TCL	Inve	rness			
	eet and No., or R.F.D. No., City, State, and Z Hubbard, OR 97032	IP Code)			FOR OFFICIAL USE ONLY PVPO NUMBER				
Characteristics of	oriate number that describes the varietal charalescribed, including numerical measurements. ANTS. Royal Horticultural Society or any rown. Descri	s, should repre ecognized cold	sent those th or fan may be	at are typical for the	se leading zeroe variety. Meas plant colors; de	es when necessary. ured data should be signate system used:			
1. SPECIF	ES: (With companion varieties for use below 1 = F. rubra ssp. commutata (Chewings)	v – use varietie 11 = Cascad 14 = Banner	e	cies of application v 12 = Highlight 15 = Barfalla	-	amestown			
	2 = F. rubra ssp. litoralis (Creeping Red)	21 = Dawson 24 = Pennlar		22 = Starlight	23 = Merlin				
	3 = F. rubra ssp. rubra (Spreading Red)	31 = Boreal 34 = Ensylva	a	32 = Ruby	33 = Fortress				
	$4 = F. \ ovina $ (Sheep)	41 = Covar							
	5 = F. longifolia (Hard)	51 = Durar		52 = Biljart (C-26	26) 53 = Scaldis				
	6 = F. tenuifolia (Fine-Leaved Sheep) 7 = Other (Specify) F	61 = Panda		62 = Barok					
2. CYTOL									
5 6	Chromosome Number 4 Ploidy	1 =	diploid	2 = tetraploid	3 = hexaploid	4 = octoploid			
. ADAPT	ATION: (0 = Not Tested; 1 = Not Adapted;	2 = Adapted)							
<u> </u>	Northeast 2 Southea Other (Specify):		2	North Central	2	Pacific N.W.			

3	Maturity Class:	ce) Location(s) of Tran(s) 22 April	2001, Hubbard, OR
	1 = Very Early (Covar)	2 = Early (Highlight)	3 = Medium Early (Boreal, Dawson)
	4 = Medium Late (Cascade, Ruby)	5 = Late (Jamestown, Agram)	6 = Very Late
	Date Headed		200400123
	Days earlier than		
· · ·	Maturity same as	Comparison Variety	
0 2	Days later than 3	4)	
5. PLAN	VT HEIGHT: (At maturity; to top of panicle; a	verage of 10 tallest culms)	
7 6 2	mm Height		
0 9 5	mm shorter than 3	4	
	Height same as	Comparison Variety	
	mm Taller than]	
6. GROV	VTH HABIT:		
2	1 = Erect (Ruby) $2 = Semi-erect (Ruby)$	Highlight) 3 = Prostrate (Si	ivana)
7. RHIZO	OMES:		
	mm Length mm Width	mm Internode length	
3	1 = Absent	2 = Weakly Creeping (Dawson)	
	3 = Strongly Creeping (Boreal)	4 = Very Strongly Creeping (Borea	al)
8. LEAF	BLADE:		
4	Color:		
	1 = Light Green (Starlight) 2 = Med	dium Light Green (Highlight)	3 = Medium Dark Green (Ruby, Agram)
* -	4 = Dark Green (Jamestown, Manoir)	5 = Bluegreen (Saphir)	6 = Graygreen (Scaldis)
	7 = Other (Specify):		
1	Glaucosity (Sowing Year): 1 = Absent	2 = Present (Vendome)	
1	Anthocycnin; $1 = Absent$ $2 = Pres$	ent Hairs (Basal):	1 = Absent 2 = Present
1	Margins: $1 = \text{Smooth}$ $2 = \text{Sem}$	i-rough 3 = Rough	
2	Margin folding (closure): 1 = Rolled inward	d (closed-Highlight) 2 = Flat	
3	Width class: 1 = Very fine (Agram, Frid	da) 2 = Fine (Jamesto	wn, Highlight, Banner, Dawson)
	3 = Medium Fine (Fortress	s, Ruby, Scaldis) 4 = Medium Coar	se (Engina)

					•							
	0 9 4	mm Length (flag leaf)										
	1 3	mm Shorter than		4		MAB 8						
		Blade length same as		$\overline{\Box}$ \gt	Comparison Varie	Z () () 4 . (00123					
		mm Longer than		ー 丿		•						
	2 4 0	mm Width (flag leaf)										
		mm Narrower than			•							
		Blade width same as			Comparison Varie	ety						
	0 2 0	mm Wider than	3	4 9								
9.	LEAF S	SHEATH:	· · · · · · · · · · · · · · · · · · ·									
		Anthocyanin (seedling):	1 = Absent (Highlight	t) $2 = Presc$	ent (Jamestown, Fo	ortress, Marga)						
	1	Auricle Hairiness:	1 = Absent	2 = Prese	ent							
	2	Margins:	1 = Open (Highlight)	1 = Open (Highlight) 2 = Closed (Jamestown)								
0.	PANIC	LE:										
- ·	1	Shape:	1 = Narrow-tapering		2 = Ovate	;						
*.			3 = Oblong		4 = Other	(Specify):						
	2	Type:	1 = Open	2 = Intern	mediate :	3 = Compact						
	1	Orientation:	1 = Erect	2 = Nodo	ling							
	2	Branch Pubescence:	1 = Glabrous	2 = Pubes	scent							
	1	Anther Color:	1 = Yellowish Green	2 = Green	ı 3 = Bluisl	ı Green	4 = Purplish					
		Glume Color	5 = Reddish	6 = Other	(Specify):							
.:	4	at 50% flowering):										
	1 1 2	mm Length										
	1 7	mm Shorter than	3	4								
		Panicle length same as			Comparison Variet	у						
		mm longer than										
ι.	PALEA:				<u> </u>							
	2	Hairs (On keels or margins	s): 1 = Absent (B	anner)	2 = Short	(Agram, Scaldis	, Olds)					
. •			3 = Long (Rai	nier, Fortress,	Jamestown)							

8.

1

LEAF BLADE (Continued):

12.	LEMM	A:			•				
1.	2	Hairs: 1 = Absent (Jamestown)	2 = Sev	veral	3 = Ma	ny (Highlight)			
. :	6 .5	mm Lemma Length					` -	1 Per Carante	····-,
	0 .6	mm Shorter than	3 4		•	20040	A 4	A **	
	•	Lemma length same as	 	Compa	arison Variety	-unda N	U I	25	
		mm Longer than							
[1 .1 0	mm Lemma Width							
[mm Narrower than							
		Lemma width same as		Compa	rison Variety				
	0 .1 0	mm Wider than	3 4						
(81	:8/4/06) 3	Awns: $1 = Absent$	2 = Present						
	1 .6	mm Awn Length							
		mm Shorter than	1						
•		Awn length same as		Compa	rison Variety				
	0 .3	mm Longer than	3 4						
13.	SEED (With lemma and palea):							
	3	Size Class (g/1000 seed):	•						
		1 = [< 09g] (Biljart, Dawson)	2 = [0.9-<].1g] (Jamestov	wn, Highlight)				
		3 = [1.1 - 1.3g] (Fortress, Novorubra)	4 = [>1.3g] (Bor	eal, Golf	rood)				
1	1 6 7	mg per 1000 seed							
	2 1 0	mg per 1000 seed less than	3 4						
		Seed Weight same as		Compai	rison Variety				
		mg per 1000 seed more than							
14.	DISEAS	E, INSECT, AND NEMATODE REACTION	Ŋ			· .			
	0	Melting-out Drechslera poae (Helminthospo	rium vagans)	0	Stripe Rust P. str	iiformis			
-	0	Leaf Spot D. siccans		6	Leaf Rust P. poac	e-nemoralis			
-	6	Net Blotch D. dictyoides		6	P. crandallii	•			
	7	Leaf Spot Bipolaris sorokiniana		0	Pythium Blight P	ythum ultimum			
: .	5	Brown Patch Rhizoctonia solani		7	Red Thread Corti	icium fusciforme			
	0	Powdery Mildew Erysiphe graminis		6	Dollar Spot Scler	otinia homoeocarp	oa -		
	5	Stripe Smut <i>Ustilago striiformis</i>			Insect				

7	F. Patch, Pink snow-mold Fusarium nivale	Nematode 2004 0 0 1 2	
6	Fusarium Blight F. trincinctum, F. roseum	Other	
0	Gray Snow Mold Typhula iotana	Other	
7	Stem Rrust Puccinia graminis	Other	

1 = Application variety is less than the comparison variety. 2 = Application variety is the same as the comparison variety.

CHARACTER	VARIETY	D.R.	CHARACTER	VARIETY	D.R.
Rhizome Length	Florentine	2	Growth Habit	Florentine	2
Leaf Width	Florentine	2	Leaf Color	Florentine	3
Panicle Color	Florentine	2	Panicle Shape	Florentine	2

Leaf W Panicle Winter Color **Florentine** Cold Injury Florentine Shade Tolerance Florentine 2 Heat **Florentine** 2 Drought Florentine 2 Disease* Florentine 3

* Specify each disease evaluated.

Michrodium patch, Leaf spot, Dollar spot, Brown patch, Pink snow mold and Summer patch

3 = More than, better, greater, darker, more disease resistant, etc.

16. ADDITIONAL DESCRIPTION:

Exhibit D.

Additional Description of Inverness (PST-47TCR, PST-47TCL) Strong Creeping Red Fescue

Inverness has good turf quality, a dark green color, good winter color, good resistance to *Michrodochium* patch, good leaf spot resistance, good dollar spot resistance, good brown patch resistance, good summer patch ratings, good red thread resistance, good pink snow mold resistance, good pink patch resistance and an improved seed yield potential (see Tables 4-22).

Table 1. 2001 mean initial heading dates for entries in a fine fescue seed yield trial seeded fall of 2000 near Hubbard, OR.

Entry	Heading Date
Bighorn	12 April
Little Bighorn	14 April
PST-4MB-98	17 April
4S3E Bulk	17 April
Seabreeze	17 April
Flyer	20 April
PST-4R3	20 April
Ensylva	20 April
Quatro	20 April
Flyer II	20 April
Shademaster	21 April
Shademaster li	21 April
Boreal	21 April
Reliant II	21 April
Inverness	22 April
Defiant	22 April
LTP-3851	22 April
Polaris	22 April
Discovery	22 April
4VB3 Bulk	23 April
LTP-HF-95-1	23 April
Aurora	23 April
PST-4BBL PST-47 TH	23 April
	23 April
LA20-0-19032	24 April
PST-4CRE-98	24 April
PST-4FRR Aurora Gold	25 April
Shadow	25 April
	25 April
Aurora II Scaldis	25 April
Florentine	25 April
Tiffany	26 April
Shadow II	27 April
PST-4FR	27 April 28 April
PST-4EC	29 April
LTP-5001	29 April
Jamestown II	29 April
Barcrown	30 April
Jamestown	03 May
Jamotovii	
LSD (0.05)	4 days

Table 1A. Mean initial heading dates for entries in a fine fescue seed yield trial seeded fall of 1998 near Hubbard, OR.

<u>Entry</u>	<u>1999</u>
4FRR	20 April
4CRE	19 April
4BP Bulk	05 May
4R3	19 April
Laxton	17 April
PST-4FR	19 April
Bighorn Flyer II	12 April 21 April
Camilla	20 April
Inverness	21 April
ZFRR-93-112X	23 April
Florentine	25 April
Syn 4V3B	22 April
Shademaster II	20 April
4UB	12 April
Syn 4S3E	16 April
Ensylva	19 April
4BBL	20 April
PST-4MB	16 April
Barnica	21 April
Syn 4UG	15 April
4BLUE	21 April
4HI	15 April
4TDD	24 April
Seabreeze Dawson	16 April
Shadow	22 April 26 April
Tiffany	28 April
4CU	23 April
Jamestown II	03 May
Shadow II	02 May
Baroxi	02 May
Syn 4RS	23 April
Discovery	22 April
Sandpiper	29 April
Aurora E	21 April
47TH	21 April
4AU	22 April
PST-4HM	22 April
Aurora Gold	22 April
Bargreen	01 May
Clio	28 April
Barcrown PST-EFL	28 April
Shademark	17 April 20 April
4PH	20 April
11 11	ZU APIII
LSD (0.05)	3 days

200400123

Table 2. 2000 mean morphological measurements for entries in a fine fescue seed yield trial seeded fall of 1998 near Hubbard, OR.

<u>Entry</u>	Plant Height <u>(cm)</u>
Dawson	90.0
Shadow	89.8
Ensylva	89.1
Seabreeze	88.4
Laxton	87.4
Shadow II	87.2
Camilla	86.0
Tiffany	86.0
47FRR	85.4
Shademaster II	84.8
Inverness	79.7
PST-EFL	79.5
PST-4S3E	78.5
4FR	76.2
Florentine	75.7
4BBL	75.6
Aurora E	73.4
Aurora Gold	70.3
Bighorn	69.6
4UB	69.1
Discovery	67.8
4BLUE	67.2
4MB	65.9
4AU	65.1
4HM	64.5
47TH	63.3
Barcrown	57.1
LSD (0.05)	2.6

Table 2A. 2001 mean morphological measurements for entries in a fine fescue seed yield trial seeded fall of 2000 near Hubbard, OR.

Entry	Plant Height (cm)
Inverness	70.1
Shademaster II	68.2
4FRR	66.6
LA20-0-19032	65.8
Flyer II	61.7
LTP-3851	60.2
LTP-5001	58.9
4CRE-98	58.1
4FR	57.4
Florentine	55.7
LTP-HF-95-1	41.8
LSD (0.05)	3.1

Table 3. 2002 mean morphological measurements for entries in a fine fescue seed yield trial seeded fall of 2000 near Hubbard, OR.

Entry	Plant Height (cm)	Panicle Length (cm)	Flag Leaf Height (cm)	Flag Leaf Length (cm)	Flag Leaf Width (mm)	Tiller Leaf Length (cm)	Tiller Leaf Width (mm)	Tiller Count (#/12.7 cm Row)
Ensylva	85.7	12.9	36.7	10.7	2.2	10.5	1.8	63.6
Inverness	76.2	11.2	25.4	9.4	2.4	8.5	1.6	96.1
LSD (0.05)	2.8	0.7	2.1	1.3	0.2	0.8	0.2	23.5

Tables 4-22 contain selected data including top and bottom commercially available entries and selected standards.

Table 4. 2000 mean turgrass quality ratings of strong creeping red fescue cultivars grown at twenty-nine locations in the US and Canada (9 = ideal)

Mean	5.8	5,8	5.7	5.7	5,6	5.5	5.	5,1	5.1	5.0	5.0	4.9	6.4	4. Q	4.8	4.7	4.6	4.3	4.2	4.1	0.2	
¥	5.8	6.4	5.7	6.1	5.5	5	5.0	5.0	4.9	5	5.0	5.2	53	4.9	4.8	4.6	3.6	4.6	4.8	5.0	0.7	
WA3	4.3	4.4	4.8	4.5	8,4	4.4	4.4	4.7	4.5	4.1	8.	4.0	4 .1	4.4	4	4.	4,9	3.5	4.0	4.0	0.8	
WA1	5.0	6.4	5.	9,	8.	4.7	4.7	5.0	5.0	5,1	6.4	4.6	4,5	4.3	4,3	4. 3.	4.	4.4	4 2	<u>4</u> .	0.5	
≶	6,3	9	5.6	5.8	9	5.8	4.7	4	5.	9,4	4	20	4.5	5.0	5,1	4.4	4	4.3	3.8	3.4	0.9	
5	4	5.2	10	4.4	4	5.0	4.4	5.0	4.9	52	0.4	4	5,5	4.5	4.2	4.0	4.3	0.4	0.4	3.6	8.0	
SD	5.3	5.3	5.8	5.7	5.6	5.3	5.4	5.3	5,4	5.6	5.3	4.3	4,5	5.1	3.8	52	5.4	4.	0.4	8.4	0.7	
æ	6.8	2	4	6.1	6.7	5.9	6.4	4.1	3.6	5.8	5.5	5.0	5.4	3.5	5.2	5.1	9	3,9	3,2	2.5	0.1	
å	5.3	'n,	5.5	4.0	5.3	5.2	5,3	5.1	5.1	5.	5.1	5.3	3.8	5.1	4.8	4.9	2.7	43	4.9	8,8	4.0	
Æ	5.8	5.6	53	5,8	5.5	53	4.4	3.9	29	3.5	3.4	5.0	5.3	2.8	6.	7.	2.7	3.9	3.0	2.7	7	
ջ	3.8	ю С	3.9	4.0	2.9	9	3,3	2.4	3.0	4	3.0	2.4	2.9	3.2	4.	2,8	 	2.0	4.	1.5	6.	
È	3.6	ю С	29	2.7	3.7	2.9	3.8	3.4	3.5	33	9.4	3.2	2.9	3.5	3.0	9.4 4.6	4.8	3.2	3.2	3.5	6.0	
Ş	8.8	6.9	6.8	6.3	8,8	6.4	4.8	5.1	0.0	5,3	5.5	6.4	2.9	5.0	5.5	4.8	3.7	4.5	4.8	4.9	2.0	
NJ2	6.1	5.9	0.0	5.5	5.5	5.7	4.3	3,3	4.	4 6	3.8	4	5.2	4 65	4 6	4.	4.4	36	3.3	3.0	8.0	
NJ1	6.4	6.5	62	5.8	5.8	5. 9.	4.9	4.6	4.6	5.0	4.8	4.6	3.9	4.8	4,8	3.8	3.9	4.	2.3	3.3	8.0	
S	5.9	0.0	5,2	6.1	6.0	8.4	4.5	4.8	53.33	9.0	8.4	5.1	5.1	0.4	5.0	4.2	5.3	3.9	4.2	3.4	1.0	
MT	2.6	5.6	2.5	5.4	9:6	5.9	5.8	5.8	5.7	6.1	5,3	5.0	5.4	5,4	5.1	5.1	3.0	5.0	5.3	5,0	7.0	
NO.	4.	20	4 60	5.0	4.	Α ω	4.3	4 .8	5.0	6.0	4.	3.7	4.3	0	3.8	3.6	6. ش	3.2	3.6	3.3	0.8	
Ξ	9.9	9	6	6.4	6.3	6.4	6.4	6.1	6.3	6.2	6.4	6.0	6.2	6.1	6.1	6.3	6.2	5.6	5.7	5.5	0.4	
ME2	12	0	7.4	6.7	6.8	6.3	8.8	7.4	6.9	2	7.	5.9	6.5	9.9	6,2	63	5.6	6.5	5.6	5.4	4.	
MET	7.2	7.7	6.9	7.0	2,5	6.7	6.8	6.4	62	7.3	6.5	2.7	6.1	6.7	5.9	6.1	53	6.1	5.6	5.9	1.0	
Đ.	4.6	9	63	9	9.9	6.4	5.7	5.8	5.4	5.7	د .	5.7	5.4	5.3	4.8	5.4	4,6	7.4	4.5	4.6	7.0	
Α̈́	5.9	S)	5,5	5.5	5,5	5.0	5.0	5.7	53	4.9	4 .8	6.4	6.4	5.2	5.0	4.9	5.7	4.4	4. 5.	4.0	7.0	
₹	4.7	2.	7.5	7.3	9.2	7.4	9.9	6.5	6.9	6.7	9.9	7.0	6.0	9.9	5.9	6.3	5.7	5,8	5.2	5.1	0.7	
ξ	7.	?	6. 4	9.2	6.0	6.9	6.0	6.0	9.0	5.4	0.0	2.0	6.1	6.0	6.1	6.3	4 .5	4.6	5.0	5.2	1.4	
3	6.6	9	7.0	8.6	6.3	5.9	0.0	20	4 .	4 . εί	5,3	5.8	4.7	8.4	5.0	1.	4	3.9	4.7	4.2	6.0	
=	5.0	ç	5,5	5.0	4.5	20	φ, 0	20	4 .0	Α. 80	4.3	5.0	4 .	4.0	4.3	4.7	4. 5.	3.8	4. 6.	4,3	7.0	
≤	7.	c)	5.4	5.4	5.	5.0	4.9	5.9	8.	7.4	5.2	7	4 8	4. 0	4.7	4.8	5.0	٠ <u>٠</u>	4.6	4.3	1.2	
₹	6.3	7	9.0	6.0	5.8	6,	5.7	6.4	6.0	5.5	6.1	5.5	6.1	6.2	5.8	5.8	5.9	5.	5.3	4.9	0.4	
¥	4.7	9	5.7	5.7	5.6	5.3	3.6	4.4	4.6	7,	4.8	3,	3.0	4.3	3.6	Ą Ĉ	4.7	3.1	3.2	3.3	1.0	
Entry	Cindy Lou	Jasper II	SRX 52961	Navigator	ABT-CR-3	PST-EFL	Inverness	PST4FR	Pathfinder	Florentine	BAR CF 8 FUS1	SR 5210	Shademaster If	DGSC 94	ASC 082	Shademark	Rose	ASC 172	Boreal	Common Creeper	LSD (0.05)	

,	ø	10010-0	
	Mean	1004001	23
	W	6.3 6.3 6.3 6.3 6.3	0.7
	WA3	5.5 5.1 5.4 4.7 7.4	9.0
	WA1	5.0 5.0 5.0 7.4 7.6 7.0 7.0 7.0 7.0 7.0 7.0 7.0 7.0 7.0 7.0	7.0
	\$	6.6. 4.4. 7.6. 7.0. 7.0. 9.0.	9.0
	5	5.6 5.0 5.0 5.0 5.0 5.0 5.0 5.0 5.0 5.0	:
	SD	8.4.4.6.6. 8.4.4.6.6. 8.6.4.6.6.6.6.6.6.6.6.6.6.6.6.6.6.6.6.6.	0.7
	2	8.3 5.8 3.1 4.2 4.2	6
	ÐE	3.4.4.4.6 6.6.6.4.0	0.3
	PA	6.6 7.7 7.2 7.3 7.4 7.5 7.7 7.7 7.7 7.7 7.7 7.7 7.7 7.7 7.7	9.0
	ĕ	4 8 4 4 8 8 8 8 9 0 0 0	0.7
Geal)	NY	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	0.7
(A = 1	SK	8.00.00 8.00.00 8.00.00	1.
Çanade	NJ2	2.2.4.4.6.9.9.9.9.9.9.9.9.9.9.9.9.9.9.9.9.9	0.6
S and	E	66.6.22.23	9.8
n me t	Š	7.50 5.60 8.40 8.50	9.0
SUOII	MT	5.1 5.9 5.9 5.7 5.7	6.0
	. Q	6.00 6.00 8.00 8.00 4.4	12
	W	0 សុសុខ្មេស 4 ១ ១ ១ សុស ប	0.7
grown at twenty-nine locations in the US and Canada (9 = 10eat	ME2	7.2 7.4 7.3 5.8 5.8	9.0
200	ME1	7.2 7.4 7.2 6.2 5.7	0.7
Cultova	Q¥ Q¥	6.8 6.8 7.0 5.3 5.3	6.4
escae	MA	0.4.4.8.8. 0.0.0.0.0.0.	6.0
8	₹	7.7 7.6 7.9 5.2 5.2	9.0
reepin	KS	7.9 7.6 7.7 7.7	6.0
	ĸ	4.3.3.4.4.8. 4.4.6.7.4.8.	0.8
32 01 5	=	5.6 5.0 5.0 5.2	0.5
	. ≰	6.3 6.3 5.6 5.6 5.8	5.
	8	5.4.4.4.4.4.4.6.6.4.4.4.4.4.4.4.4.4.4.4.	0.2
Turgrass	Ϋ́	5.9 5.5 6.4 5.6 5.4 7.9 7.7 5.6 7.4 7.2 7.7 5.6 4.7 6.3 4.9 6.2 7.8 7.6 7.0 4.5 6.8 7.1 7.5 6.3 4.6 6.1 5.0 4.7 7.7 7.3 4.3 6.7 7.2 7.9 7.7 7.3 4.3 6.7 7.2 7.9 7.9 7.9 4.3 6.7 7.7 7.3 4.3 6.7 7.2 7.9 7.9 4.4 5.6 5.0 4.4 5.4 5.4 5.2 3.9 5.1 6.2 5.9 5.1 3.7 5.3 5.7 5.9 5.1 5.1 5.7 5.3 5.7 5.9 5.1 5.1 5.1 5.2 5.7 5.9 5.1 5.1 5.2 5.3 5.7 5.9 5.1 5.1 5.2 5.3 5.7 5.9 5.1 5.1 5.2 5.3 5.7 5.9 5.1 5.1 5.2 5.3 5.7 5.9 5.1 5.2 5.3 5.7 5.9 5.1 5.2 5.3 5.7 5.9 5.2 5.3 5.7 5.9 5.2 5.3 5.7 5.9 5.2 5.3 5.7 5.9 5.2 5.3 5.7 5.9 5.2 5.3 5.7 5.9 5.2 5.3 5.7 5.9 5.2 5.9 5.7 5.9 5.2 5.9 5.1 5.2 5.9 5.7 5.9 5.9 5.7 5.9 5.9 5.7 5.9 5.9 5.7 5.9 5.9 5.7 5.9 5.9 5.7 5.9 5.9 5.7 5.9 5.9 5.7 5.9 5.9 5.7 5.9 5.9 5.7 5.9 5.9 5.7 5.9 5.9 5.7 5.9 5.9 5.7 5.9 5.9 5.7 5.9 5.9 5.7 5.9 5.9 5.7 5.9 5.9 5.9 5.9 5.9 5.9 5.9 5.9 5.9 5.9	0.7
table 5. 1999 mean thirthrass qualify ratings of strong creeping red rescue cultivars.	Entry	Jasper II Florentine Shademaster II Inverness Boreal Common Creeper	LSD (0.05)

200400123

7.2 7.0 6.9 6.9 5.5

880 7.7.0 6.0 6.3

8.0 7.7 7.3 7.0 7.0

7.7 6.0 6.0 6.0 5.3

6.3 7.7 6.3 4.7 5.3 6.3 6.3 7.7 6.3 6.7 5.0 6.0 6.0 6.7 6.0 6.3 6.0 6.0 6.0 7.0 6.0 6.3 6.0 7.0 5.0 3.3 5.3 5.7 6.0 4.3 3.7 5.3

7.3 9.0 7.0 8.0 7.3 8.0 6.7 7.0 6.0 5.7 5.7 4.7

7.0 7.0 7.0 6.0 5.3

7.0 7.7 7.3 7.3 6.3 7.3 6.0 6.3 5.7 6.3 6.3 5.0

9.0 9.0 7.3 7.3

8.0 7.3 6.7 6.3 5.7

6.0 7.0 5.7 7.0 6.7 6.3 6.7 6.3 5.3 6.0 4.7 5.3

8.3 7.7 7.3 8.0 6.3

8.7 7.7 7.3 5.3 5.3

9.0 7.3 6.0 7.0 7.0 7.0

5.3 5.3 5.3 5.3

8.0 7.0 7.0 7.0 8.0 7.0 7.3 6.3 6.3 5.7 8.9 6.3

6.0 6.3 6.3 8.3 8.3 8.3 4.

6.0

7

6.0

9.

7

970

Mean	7.2 7.0 7.0 6.9 6.4 6.1	0.2
WIZ	7.0 7.0 7.0 6.7 6.3	0.7
WII	6.7 6.7 6.7 6.7	9.6
WA3	8.0 7.3 7.0 6.7 6.0 6.0	0.6
WA1	7.0 7.7 7.7 7.3 6.3 5.7	0,10
¥	6.7 6.7 6.7 6.0 6.0	2
5	8 4 8 8 4 4 6 6 6 6 6	4.
Ŝ	7.7 7.3 7.0 5.7 5.3	6.0
묎	8.3 7.7 6.7 6.0 6.0	12
쁑	7.0 7.0 7.0 7.0 5.3	0.7
ş	7.3 7.3 7.0 7.0 7.0	9.6
¥	7.0 7.0 7.0 6.0 6.3	1.5
SN	7.7 6.0 6.0 6.3	1.7 dark g
S	8.0 6.7 3.7 5.0 5.0	1.6 a (9 =
뵕	5.0 7.7 5.3 5.3	1.4 Canad
NC2	8.7 7.3 6.7 8.0 8.0	1.5 1.4 1.6 US and Canada (9 = c
NC	7.7 8.0 8.0 6.3	1.6 sin the 0
M	8.8.2 8.3.3 8.3.5	1.3
M12	0.2 0.2 0.2 0.2 0.2 0.2 0.2	0.5 -four to
MIT	7.7 7.0 7.0 7.0 7.0 7.0	0.2 twent
QW	6.3 6.3 6.3	0.7 оwn at
≨	6.7 6.0 6.0 7.3 6.7	2.1
₹	9.0 9.0 9.0 7.7 8.0	0.7 e cuttin
KS	7.7 8.7 7.3 7.0 6.7	1.0 fescu
ĸ	6.7 6.7 4.7 5.0	1.1 1.2 1.0 0.7 creeping red fescue cutti
교	844.004.4 00.000	- Gee
AR CA IA IL IN KS KY	4.00.00.00 6.00.00 7.00 7.00	1.2 strong o
8	6.7 6.3 6.3 3.7	1.6 ngs of s
AR	8.7 7.0 7.3 6.0 6.0	1.2 olor rating
Entry	ASC 172 Inverness Florentine Shademaster It Boreal Common Creeper	LSD (0.05) 1.2 1.6 1.2 1.1 1.2 1.0 0.7 2.1 0.7 0.2 0.1 Table 7. 1999 genetic color ratings of strong creeping red fescue cultivars grown at twenty-four

Table 8. 2000 mean winter color ratings of strong creeping red fescue cultivars grown at 2 locations in the US (9 = complete color retention)

Entry	OK1	VA1	Mean		
Common Crooning Rad	6.0	5.0	5.5	20040012	7
Common Creeping Red Florentine	5.7	5.3	5.5 5.5	= 0 0 1 0 1 E	Ų)
Shademaster II	5.0	6.0	5.5		
Boreal	6.0	4.7	5 .3		
Inverness	5.0	4.3	4.7		
SRX 52961	5.0	4.0	4.5		
LSD (0.05)	0.7	1.1	0.7		

Table 9. 1999 mean winter color ratings of strong creeping red fescue cultivars grown at 4 locations in the US (9 = complete color retention)

Entry	KY1	NJ1	OK1	VA1	Mean
Inverness	7.0	7.0	4.3	6.7	6.3
Florentine	7.7	5.3	4.3	6.3	5.9
Shademaster if	6.7	6.0	4.3	5.7	5.7
Boreal	5.3	5.0	3.7	4.7	4.7

Table 10. 1999 mean Microdochium patch ratings of strong creeping red fescue cultivars grown at 2 locations in the US (9 = no disease)

Entry	ME1	ME2	Mean
Salsa	8.0	8.3	8.2
	7.7	8.3	8.0
Shademaster II			
Florentine	7.7	7.0	7.3
Boreal	7.7	6.3	7.0
Inverness	7.3	6.3	6.8
Jasper II	7.3	6.3	6.8
LSD (0.05)	2.2	1.8	1.4

Table 11. 2000 mean leaf spot ratings of strong creeping red fescue cultivars grown at 1 location in the US (9 = no disease)

Entry	NJ2
leener II	5.7
Jasper II Florentine	4.7
Inverness	4.7
Shademaster II	3.0
Boreal	2.0
LSD (0.05)	1.3

Table 12. 1999 mean leaf spot ratings of strong creeping red fescue cultivars grown at 4 locations in the US (9 = no disease)

		-			
Entry	ME1	ME2	NJ1	NJ2	Mean
Jasper II	5.7	8.0	4.0	5.7	5.8
Inverness	5.3	7.3	2.7	5.7	5.3
Florentine	5.3	8.0	2.0	4.3	4.9
Shademaster II	5.7	6.0	2.7	4.3	4.7
Boreal	3.3	4.3	2.7	2.0	3.1
Shademark	3.0	4.7	2.0	2.3	3.0
LSD (0.05)	1.9	4.0	0.9	1.4	1.2

Table 13. 2000 mean dollar spot ratings of strong creeping red fescue cultivars grown at 4 locations in the US (9 = no disease)

Entry	IN1	NJ2	PA1	WI2	Mean	
Jasper II	7.7	8.0	7.7	5.0	7.1	
Inverness	6.7	4.0	7.0	4.7	5.6	200400123
Boreal	6.3	3.0	5.7	4.0	4.8	
Florentine	6.0	3.0	4.7	4.0	4.4	
Shademaster II	7.3	2.7	3.3	4.3	4.4	
Pathfinder	5.3	1.0	3.0	3.3	3.2	
LSD (0.05)	2.5	1.7	2.3	0.9	1.0	

Table 14. 1999 mean dollar spot ratings of strong creeping red fescue cultivars grown at 1 location in the US (9 = no disease)

Entry	PA1
Inverness	6.3
Florentine	6.0
Shademaster II	5.7
Boreal	4.3
LSD (0.05)	1.7

Table 15. 1999 mean brown patch ratings of strong creeping red fescue cultivars grown at 2 locations in the US (9 = no disease)

Entry	ME1	ME2	Mean
Florentine	7.7	6.7	7.2
Shademaster II	7.3	5.3	6.3
Inverness	5.7	6.0	5.8
Boreal	3.7	3.7	3.7
Common Creeping Red	3.3	2.7	3.0
LSD (0.05)	2.3	2.6	1.7

Table 16. 2000 mean summer patch ratings of strong creeping red fescue cultivars grown at 2 locations in the US (9 = no disease)

Entry	NJ1	Wi2	Mean
Jasper II	7.7	8.3	8.0
Shademaster II	6.7	6.7	6.7
Florentine	5.7	6.3	6.0
Inverness	4.3	7.0	5.7
Boreal	2.0	6.3	4.2
LSD (0.05)	2.2	1.8	1.4

Table 17. 1999 mean summer patch ratings of strong creeping red fescue cultivars grown at 1 location in the US (9 = no disease)

Entry	NC1
Jasper II	8.0
Florentine	7.7
Inverness	7.3
Shademaster II	7.3
Boreal	6.0
Common Creeping Red	5.0
LSD (0.05)	1.3

Table 18. 2000 mean red thread ratings of strong creeping red fescue cultivars grown at 4 locations in the US and Canada (9 = no disease)

Entry	ME2	NJ2	NS1	WA3	Mean	
Boreal	7.3	8.0	5.7	6.0	6.8	200400123
Inverness	8.7	7.3	5.3	5.3	6.7	
Florentine	9.0	6.3	4.3	5.7	6.3	
Shademaster II	8.7	5.3	5.3	6.0	6.3	•
LSD (0.05)	2.5	3.4	2.2	1.6	1.3	

Table 19. 2000 mean pink snow mold ratings of strong creeping red fescue cultivars grown at 1 location in the US (9 = no disease)

Entry	ME2
D-0-6-1	
Pathfinder	8.7
Florentine	8.0
Inverness	8.0
Shademaster II	6.7
Boreal	6.0
Shademark	5.3
LSD (0.05)	1.7

Table 20. 2000 mean red thread/pink patch ratings of strong creeping red fescue cultivars grown at 1 location in the US (9 = no disease)

Entry	PA1
Shademaster II	5.7
Inverness	5.3
Florentine	5.0
Boreal	4.7
LSD (0.05)	1.3

Table 21. Mean turf quality and leaf spot ratings for strong creeping red fescue entries in a fine fescue turf trial seeded fall of 1998 near Hubbard, OR. (9 = ideal; no disease)

	T	Leaf Spot		
Entry	1999	2000	Mean	1999
Florentine	5.4	5.3	5.3	6.3
Shademaster II	5.2	5.0	5.1	6.3
Inverness	6.3	5.1	5.7	5.3
Boreal	4.6	4.8	4.7	4.7
LSD (0.05)	0.7	0.8	0.6	1.3

Table 22..2000 mean seed yield for entries in a fine fescue seed yield trial seeded fall of 1998 near Hubbard, OR.

Entry	1999 (lb/A)	2000 (lb/A)	Mean (lb/A)
Inverness	1365.2	1378.4	1397.5
Shademaster II	1525.3	1000.7	1263
Flyer II	756,3	862.4	809.3

REPRODUCE LOCALLY. Include form number and edition date on all	reproductions. F	ORM APPROVED - OMB No. 0581-0055	
U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHIP	Application is required in order to detectificate is to be issued (7 U.S.C. 24 confidential until the certificate is issued	21). The information is held	
1. NAME OF APPLICANT(S)	2. TEMPORARY DESIGNATION	3. VARIETY NAME	
• •	OR EXPERIMENTAL NUMBER	3. VARIETY NAME	
Pure Seed Testing, Inc.	PST-47TCR, PST-47TCL	Inverness	
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5. TELEPHONE (Include area code) 6. FAX (Include area code)		
P.O. Box 449	(503) 651-2130	(503) 263-0703	
Hubbard, OR 97032	7. PVPO NUMBER		
		200400123	
8. Does the applicant own all rights to the variety? Mark an "X" in the	t e appropriate block. If no, please explai	n. YES NO	
9. Is the applicant (individual or company) a U.S. national or a U.S. ba	ased company? If no, give name of co	ountry. YES NO	
10. Is the applicant the original owner?	NO If no, please answer <u>one</u> o	of the following:	
a. If the original rights to variety were owned by individual(s), is (a	no If no, give name of country		
b. If the original rights to variety were owned by a company(ies),	is (are) the original owner(s) a U.S. bas		
11. Additional explanation on ownership (Trace ownership from origin Pure Seed Testing has granted Turf-Seed, Inc. an exclusive license throught the world.			
PLEASE NOTE:			
Plant variety protection can only be afforded to the owners (not license	ees) who meet the following criteria:		
If the rights to the variety are owned by the original breeder, that penational of a country which affords similar protection to nationals of	rson must be a U.S. national, national o the U.S. for the same genus and specie	f a UPOV member country, or ss.	
If the rights to the variety are owned by the company which employed nationals of a UPOV member country, or owned by nationals of a co- genus and species.	ed the original breeder(s), the company ountry which affords similar protection to	must be U.S. based, owned by nationals of the U.S. for the same	
3. If the applicant is an owner who is not the original owner, both the o	riginal owner and the applicant must me	eet one of the above criteria.	
The original breeder/owner may be the individual or company who direct for definitions.	ected the final breeding. See Section 4	(a)(2) of the Plant Variety Protection	
According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, a control number. The valid OMB control number for this information collection is 0581-0055. I including the time for reviewing the instructions, searching existing data sources, gathering an	The time required to complete this information collect	on is actimated to everage R 1 hour per reconnec	
The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and act marital or family status, political beliefs, parentel status, or protected genetic information. (Not communication of program information (Braille, large print, audiotape, etc.) should contact US.	tivities on the basis of race, color, national origin, gen	der, religion, age, disability, sexual orientation,	

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th end Independence Avenue, SW, Washington, D.C. 20250-9410 or cell (202) 720-5964 (voice and TDD). USDA is an equal opportunity provide and employer.

ST-470-E (04-03) designed by the Plant Variety Protection Office using Word 2000